



On Solé and Planat Criterion for the Riemann Hypothesis

Frank Vega

EasyChair preprints are intended for rapid dissemination of research results and are integrated with the rest of EasyChair.

September 23, 2023

On Solé and Planat criterion for the Riemann Hypothesis

Frank Vega

NataSquad, 10 rue de la Paix 75002 Paris, France

Abstract

The Riemann hypothesis is the assertion that all non-trivial zeros have real part $\frac{1}{2}$. It is considered by many to be the most important unsolved problem in pure mathematics. There are several statements equivalent to the famous Riemann hypothesis. In 2011, Solé and Planat stated that the Riemann hypothesis is true if and only if the inequality $\zeta(2) \cdot \prod_{q \leq q_n} (1 + \frac{1}{q}) > e^\gamma \cdot \log \theta(q_n)$ holds for all prime numbers $q_n > 3$, where $\theta(x)$ is the Chebyshev function, $\gamma \approx 0.57721$ is the Euler-Mascheroni constant, $\zeta(x)$ is the Riemann zeta function and \log is the natural logarithm. In this note, using Solé and Planat criterion, we prove that the Riemann hypothesis is true.

Keywords: Riemann hypothesis, Riemann zeta function, Prime numbers, Chebyshev function
2000 MSC: 11M26, 11A41, 11A25

1. Introduction

In mathematics, the Riemann hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part $\frac{1}{2}$. It was proposed by Bernhard Riemann (1859). The Riemann hypothesis belongs to the Hilbert's eighth problem on David Hilbert's list of twenty-three unsolved problems. This is one of the Clay Mathematics Institute's Millennium Prize Problems.

In number theory, the Chebyshev function $\theta(x)$ is given by

$$\theta(x) = \sum_{q \leq x} \log q$$

with the sum extending over all prime numbers q that are less than or equal to x , where \log is the natural logarithm.

Proposition 1.1. *By proving the existence of a zero-free region for the Riemann zeta function, de la Vallée-Poussin was able to prove that [1]:*

$$\theta(x) = x + O(x \cdot \exp(-c_2 \cdot \sqrt{\log x}))$$

where c_2 is a positive absolute constant.

Leonhard Euler studied the following value of the Riemann zeta function (1734).

Email address: vega.frank@gmail.com (Frank Vega)

Preprint submitted to Elsevier

September 22, 2023

Proposition 1.2. *It is known that [2, (1) pp. 1070]:*

$$\zeta(2) = \prod_{k=1}^{\infty} \frac{q_k^2}{q_k^2 - 1} = \frac{\pi^2}{6},$$

where q_k is the k th prime number (We also use the notation q_n to denote the n th prime number).

Franz Mertens obtained some important results about the constants B and H (1874). We define $H = \gamma - B$ such that $B \approx 0.2614972128$ is the Meissel-Mertens constant and $\gamma \approx 0.57721$ is the Euler-Mascheroni constant [3, (17.) pp. 54]. On the sum of the reciprocals of all prime numbers not exceeding x , we have

Proposition 1.3. *In 1909 Edmund Landau, by using the best version of the prime number theorem then at his disposition, proved that [4]:*

$$\sum_{q \leq x} \frac{1}{q} = \log \log x + B + O(\exp(-\sqrt[4]{\log x})).$$

Proposition 1.4. *There are infinitely many natural numbers x such that [5]:*

$$B + \log \log(x) > \sum_{q \leq x} \frac{1}{q}.$$

In mathematics, $\Psi(n) = n \cdot \prod_{q|n} \left(1 + \frac{1}{q}\right)$ is called the Dedekind Ψ function, where $q | n$ means the prime q divides n . We say that $\text{Dedekind}(q_n)$ holds provided that

$$\prod_{q \leq q_n} \left(1 + \frac{1}{q}\right) > \frac{e^\gamma}{\zeta(2)} \cdot \log \theta(q_n).$$

Next, we have Solé and Planat Theorem:

Proposition 1.5. *Dedekind(q_n) holds for all prime numbers $q_n > 3$ if and only if the Riemann hypothesis is true [6, Theorem 4.2 pp. 5].*

A natural number N_k is called a primorial number of order k precisely when,

$$N_k = \prod_{i=1}^k q_i.$$

We define $R(n) = \frac{\Psi(n)}{n \cdot \log \log n}$ for $n \geq 3$. $\text{Dedekind}(q_n)$ holds if and only if $R(N_n) > \frac{e^\gamma}{\zeta(2)}$ is satisfied. There are several statements out from the Riemann hypothesis assumption:

Proposition 1.6. *We have [6, Proposition 3. pp. 3]:*

$$\lim_{k \rightarrow \infty} R(N_k) = \frac{e^\gamma}{\zeta(2)}.$$

The following property is based on natural logarithms:

Proposition 1.7. *For $x > -1$ [7, pp. 1]:*

$$\frac{x}{x+1} \leq \log(1+x) \leq x.$$

Putting all together yields a proof for the Riemann hypothesis using the Chebyshev function.

2. What if the Riemann hypothesis were false?

Several analogues of the Riemann hypothesis have already been proved. Many authors expect (or at least hope) that it is true. However, there are some implications in case of the Riemann hypothesis might be false.

Lemma 2.1. *If the Riemann hypothesis is false, then there are infinitely many prime numbers q_n for which $\text{Dedekind}(q_n)$ fails (i.e. $\text{Dedekind}(q_n)$ does not hold).*

Proof. The Riemann hypothesis is false, if there exists some natural number $x_0 \geq 5$ such that $g(x_0) > 1$ or equivalent $\log g(x_0) > 0$:

$$g(x) = \frac{e^\gamma}{\zeta(2)} \cdot \log \theta(x) \cdot \prod_{q \leq x} \left(1 + \frac{1}{q}\right)^{-1}.$$

We know the bound [6, Theorem 4.2 pp. 5]:

$$\log g(x) \geq \log f(x) - \frac{2}{x}$$

where f was introduced in the Nicolas paper [8, Theorem 3 pp. 376]:

$$f(x) = e^\gamma \cdot \log \theta(x) \cdot \prod_{q \leq x} \left(1 - \frac{1}{q}\right).$$

When the Riemann hypothesis is false, then there exists a real number $b < \frac{1}{2}$ for which there are infinitely many natural numbers x such that $\log f(x) = \Omega_+(x^{-b})$ [8, Theorem 3 (c) pp. 376]. According to the Hardy and Littlewood definition, this would mean that

$$\exists k > 0, \forall y_0 \in \mathbb{N}, \exists y \in \mathbb{N} (y > y_0): \log f(y) \geq k \cdot y^{-b}.$$

That inequality is equivalent to $\log f(y) \geq \left(k \cdot y^{-b} \cdot \sqrt{y}\right) \cdot \frac{1}{\sqrt{y}}$, but we note that

$$\lim_{y \rightarrow \infty} \left(k \cdot y^{-b} \cdot \sqrt{y}\right) = \infty$$

for every possible positive value of k when $b < \frac{1}{2}$. In this way, this implies that

$$\forall y_0 \in \mathbb{N}, \exists y \in \mathbb{N} (y > y_0): \log f(y) \geq \frac{1}{\sqrt{y}}.$$

Hence, if the Riemann hypothesis is false, then there are infinitely many natural numbers x such that $\log f(x) \geq \frac{1}{\sqrt{x}}$. Since $\frac{2}{x} = o\left(\frac{1}{\sqrt{x}}\right)$, then it would be infinitely many natural numbers x_0 such that $\log g(x_0) > 0$. In addition, if $\log g(x_0) > 0$ for some natural number $x_0 \geq 5$, then $\log g(x_0) = \log g(q_n)$ where q_n is the greatest prime number such that $q_n \leq x_0$. Actually,

$$\prod_{q \leq x_0} \left(1 + \frac{1}{q}\right)^{-1} = \prod_{q \leq q_n} \left(1 + \frac{1}{q}\right)^{-1}$$

and

$$\theta(x_0) = \theta(q_n)$$

according to the definition of the Chebyshev function. □

3. Main Insight

This is the main insight.

Theorem 3.1. *The Riemann hypothesis is true when for every large enough prime number $q_n > 3$, there exists another prime $q_{n'} > q_n$ such that*

$$R(N_{n'}) \leq R(N_n).$$

Proof. If the Riemann hypothesis is false and the inequality

$$R(N_{n'}) \leq R(N_n)$$

is satisfied for every large enough prime number $q_n > 3$, then there is an infinite subsequence of natural numbers n_i such that

$$R(N_{n_{i+1}}) \leq R(N_{n_i}),$$

$q_{n_{i+1}} > q_{n_i}$ and $\text{Dedekind}(q_{n_i})$ fails by Lemma 2.1.

This is a contradiction with the fact that

$$\liminf_{k \rightarrow \infty} R(N_k) = \lim_{k \rightarrow \infty} R(N_k) = \frac{e^\gamma}{\zeta(2)}$$

by Proposition 1.6. By definition of the limit inferior for any positive real number ε , only a finite number of elements of the sequence $R(N_k)$ are less than $\frac{e^\gamma}{\zeta(2)} - \varepsilon$. This is a contradiction with the existence of previous infinite subsequence and thus, the Riemann hypothesis must be true. \square

4. Main Theorem

This is the main theorem.

Theorem 4.1. *The Riemann hypothesis is true.*

Proof. The Riemann hypothesis is true when

$$R(N_{n'}) \leq R(N_n)$$

is satisfied for large enough prime numbers $q_{n'} > q_n$ because of the Theorem 3.1. That is the same as

$$\frac{\prod_{q \leq q_{n'}} \left(1 + \frac{1}{q}\right)}{\log \theta(q_{n'})} \leq \frac{\prod_{q \leq q_n} \left(1 + \frac{1}{q}\right)}{\log \theta(q_n)}$$

which is

$$\prod_{q_n < q \leq q_{n'}} \left(1 + \frac{1}{q}\right) \leq \frac{\log \theta(q_{n'})}{\log \theta(q_n)}$$

and finally,

$$\log \log \theta(q_{n'}) - \log \log \theta(q_n) \geq \sum_{q_n < q \leq q_{n'}} \log \left(1 + \frac{1}{q}\right)$$

after making a simple distribution and applying the logarithm to the inequality. By Propositions 1.1, 1.3 and 1.7, we have for every large enough prime number $q_n > 3$, there always exists another prime $q_{n'} > q_n$ such that

$$\begin{aligned}
& \log \log \theta(q_{n'}) - \log \log \theta(q_n) \\
&= \log \log \theta(x) - \log \log \theta(q_n) \\
&\geq \log \log \left(x + O(x \cdot \exp(-c_2 \cdot \sqrt{\log x})) \right) - \log \log \left(q_n + O(q_n \cdot \exp(-c_2 \cdot \sqrt{\log q_n})) \right) \\
&= \log \left((\log x) \cdot \left(1 + \frac{\log \left(1 + O(\exp(-c_2 \cdot \sqrt{\log x})) \right)}{\log x} \right) \right) \\
&\quad - \log \left((\log q_n) \cdot \left(1 + \frac{\log \left(1 + O(\exp(-c_2 \cdot \sqrt{\log q_n})) \right)}{\log q_n} \right) \right) \\
&= \log \log(x) - \log \log(q_n) \\
&\quad + \log \left(1 + \frac{\log \left(1 + O(\exp(-c_2 \cdot \sqrt{\log x})) \right)}{\log x} \right) - \log \left(1 + \frac{\log \left(1 + O(\exp(-c_2 \cdot \sqrt{\log q_n})) \right)}{\log q_n} \right) \\
&\geq \log \log(x) - \log \log(q_n) \\
&\quad + \frac{O(\exp(-c_2 \cdot \sqrt{\log x}))}{(\log x) \cdot (O(\exp(-c_2 \cdot \sqrt{\log x})) + 1) + O(\exp(-c_2 \cdot \sqrt{\log x}))} - \frac{O(\exp(-c_2 \cdot \sqrt{\log q_n}))}{\log q_n} \\
&\geq \log \log(x) - \log \log(q_n) - O(\exp(-\sqrt[4]{\log q_n})) \\
&\geq \log \log(x) + B - \sum_{q \leq q_n} \frac{1}{q} \\
&\geq \sum_{q_n < q \leq q_{n'}} \frac{1}{q} \\
&\geq \sum_{q_n < q \leq q_{n'}} \log \left(1 + \frac{1}{q} \right)
\end{aligned}$$

where

$$B + \log \log(x) > \sum_{q \leq x} \frac{1}{q} = \sum_{q \leq q_{n'}} \frac{1}{q}$$

holds by Proposition 1.4 when $\theta(q_{n'}) = \theta(x)$. Consequently, the inequality

$$R(N_{n'}) \leq R(N_n)$$

holds for sufficiently large prime numbers $q_{n'} > q_n$ and therefore, the Riemann hypothesis is true. \square

Acknowledgments

The author thanks Emmanuel (CEO of NataSquad) for the financial support.

References

- [1] C. J. d. L. V. Poussin, Sur la fonction $\zeta(s)$ de Riemann et le nombre des nombres premiers inférieurs à une limite donnée, Vol. 51, Hayez, 1899.
- [2] R. Ayoub, Euler and the zeta function, *The American Mathematical Monthly* 81 (10) (1974) 1067–1086. doi: 10.2307/2319041.
- [3] F. Mertens, Ein Beitrag zur analytischen Zahlentheorie., *J. reine angew. Math.* 1874 (78) (1874) 46–62. doi: 10.1515/crll.1874.78.46.
- [4] E. Landau, *Handbuch der Lehre von der Verteilung der Primzahlen*, Teubner, 1909.
- [5] G. Robin, Sur l'ordre maximum de la fonction somme des diviseurs, *Séminaire Delange-Pisot-Poitou, Théorie des nombres (1981-1982)*. *Progress in Mathematics* 38 (1983) 233–244.
- [6] P. Solé, M. Planat, Extreme values of the Dedekind ψ function, *Journal of Combinatorics and Number Theory* 3 (1) (2011) 33–38.
- [7] L. Kozma, Useful Inequalities, http://www.lkozma.net/inequalities_cheat_sheet/ineq.pdf, Accessed 22 September 2023 (2023).
- [8] J.-L. Nicolas, Petites valeurs de la fonction d'Euler, *Journal of number theory* 17 (3) (1983) 375–388. doi:10.1016/0022-314X(83)90055-0.