



Properties of the Robin's Inequality

Frank Vega

EasyChair preprints are intended for rapid dissemination of research results and are integrated with the rest of EasyChair.

September 13, 2020

PROPERTIES OF THE ROBIN'S INEQUALITY

FRANK VEGA

ABSTRACT. The Riemann hypothesis is considered the most important unsolved problem in mathematics. The Robin's inequality is true for every natural number $n > 5040$ if and only if the Riemann hypothesis is true. We prove the Robin's inequality is true for every natural number $n > 5040$ when n is not divisible by any prime number $q_m \leq 113$. In addition, the Robin's inequality is true for every natural number $n = 113^k \times n' > 5040$ when $(\ln n')^\beta \leq \ln n$ such that $\beta = \frac{113}{112}$, $113 \nmid n'$ and for an integer $k \geq 1$. Moreover, given a natural number $n = q_1^{a_1} \times q_2^{a_2} \times \cdots \times q_m^{a_m}$ such that $n > 5040$, q_1, q_2, \dots, q_m are prime numbers and a_1, a_2, \dots, a_m are positive integers, then the Robin's inequality is true for n when $q_1^\alpha \times q_2^\alpha \times \cdots \times q_m^\alpha \leq n$, where $\alpha = (\ln n')^\beta$, $\beta = (\frac{\pi^2}{6} - 1)$ and n' is the squarefree kernel of n .

1. INTRODUCTION

In mathematics, the Riemann hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part $\frac{1}{2}$. Many consider it to be the most important unsolved problem in pure mathematics [2]. It is of great interest in number theory because it implies results about the distribution of prime numbers [2]. It was proposed by Bernhard Riemann (1859), after whom it is named [2]. It is one of the seven Millennium Prize Problems selected by the Clay Mathematics Institute to carry a US 1,000,000 prize for the first correct solution [2].

The divisor function $\sigma(n)$ for a natural number n is defined as the sum of the powers of the divisors of n

$$\sigma(n) = \sum_{k|n} k$$

where $k | n$ means that the natural number k divides n [6]. In 1915, Ramanujan proved that under the assumption of the Riemann hypothesis, the inequality

$$\sigma(n) < e^\gamma \times n \times \ln \ln n$$

holds for all sufficiently large n , where $\gamma \approx 0.57721$ is the Euler-Mascheroni constant [2]. The largest known value that violates the inequality is $n = 5040$. In 1984, Guy Robin proved that the inequality is true for all $n > 5040$ if and only if the Riemann hypothesis is true [2]. Using this inequality, we show an interesting result.

2. RESULTS

Theorem 2.1. *Given a natural number*

$$n = q_1^{a_1} \times q_2^{a_2} \times \cdots \times q_m^{a_m}$$

2010 *Mathematics Subject Classification.* Primary 11M26; Secondary 11A41.
Key words and phrases. number theory, inequality, divisor, prime.

such that q_1, q_2, \dots, q_m are prime numbers and a_1, a_2, \dots, a_m are positive integers, then we obtain the following inequality

$$\frac{\sigma(n)}{n} < \frac{\pi^2}{6} \times \prod_{i=1}^m \frac{q_i + 1}{q_i}.$$

Proof. From the article reference [1], we know that

$$(2.1) \quad \frac{\sigma(n)}{n} < \prod_{i=1}^m \frac{q_i}{q_i - 1}.$$

We can easily prove that

$$\prod_{i=1}^m \frac{q_i}{q_i - 1} = \prod_{i=1}^m \frac{1}{1 - q_i^{-2}} \times \prod_{i=1}^m \frac{q_i + 1}{q_i}.$$

However, we know that

$$\prod_{i=1}^m \frac{1}{1 - q_i^{-2}} < \prod_{j=1}^{\infty} \frac{1}{1 - q_j^{-2}}$$

where q_j is the j^{th} prime number and

$$\prod_{j=1}^{\infty} \frac{1}{1 - q_j^{-2}} = \frac{\pi^2}{6}$$

as a consequence of the result in the Basel problem [6]. Consequently, we obtain that

$$\prod_{i=1}^m \frac{q_i}{q_i - 1} < \frac{\pi^2}{6} \times \prod_{i=1}^m \frac{q_i + 1}{q_i}$$

and thus,

$$\frac{\sigma(n)}{n} < \frac{\pi^2}{6} \times \prod_{i=1}^m \frac{q_i + 1}{q_i}.$$

□

Theorem 2.2. For $x \geq 11$, we have

$$\sum_{q \leq x} \frac{1}{q} < \ln \ln x + \gamma - 0.12$$

where $q \leq x$ means all the primes lesser than or equal to x .

Proof. For $x > 1$, we have

$$\sum_{q \leq x} \frac{1}{q} < \ln \ln x + B + \frac{1}{\ln^2 x}$$

where

$$B = 0.2614972128 \dots$$

is the (Meissel-)Mertens constant, since this is a proven result from the article reference [4]. This is the same as

$$\sum_{q \leq x} \frac{1}{q} < \ln \ln x + \gamma - \left(C - \frac{1}{\ln^2 x}\right)$$

where $\gamma - B = C > 0.31$, because of $\gamma > B$. If we analyze $(C - \frac{1}{\ln^2 x})$, then this complies with

$$(C - \frac{1}{\ln^2 x}) > (0.31 - \frac{1}{\ln^2 11}) > 0.12$$

for $x \geq 11$ and thus, we finally prove that

$$\sum_{q \leq x} \frac{1}{q} < \ln \ln x + \gamma - (C - \frac{1}{\ln^2 x}) < \ln \ln x + \gamma - 0.12.$$

□

Definition 2.3. We recall that an integer n is said to be squarefree if for every prime divisor q of n we have $q^2 \nmid n$, where $q^2 \nmid n$ means that q^2 does not divide n [1].

Theorem 2.4. *Given a squarefree number*

$$n = q_1 \times \cdots \times q_m$$

such that q_1, q_2, \dots, q_m are odd prime numbers, the greatest prime divisor of n is greater than 7 and $3 \nmid n$, then we obtain the following inequality

$$\frac{\pi^2}{6} \times \frac{3}{2} \times \sigma(n) \leq e^\gamma \times n \times \ln \ln(2^{19} \times n).$$

Proof. This proof is very similar with the demonstration in Theorem 1.1 from the article reference [1]. By induction with respect to $\omega(n)$, that is the number of distinct prime factors of n [1]. Put $\omega(n) = m$ [1]. We need to prove the assertion for those integers with $m = 1$. From a squarefree number n , we obtain that

$$(2.2) \quad \sigma(n) = (q_1 + 1) \times (q_2 + 1) \times \cdots \times (q_m + 1)$$

when $n = q_1 \times q_2 \times \cdots \times q_m$ [1]. In this way, for every prime number $q_i \geq 11$, then we need to prove that

$$(2.3) \quad \frac{\pi^2}{6} \times \frac{3}{2} \times (1 + \frac{1}{q_i}) \leq e^\gamma \times \ln \ln(2^{19} \times q_i).$$

For $q_i = 11$, we have that

$$\frac{\pi^2}{6} \times \frac{3}{2} \times (1 + \frac{1}{11}) \leq e^\gamma \times \ln \ln(2^{19} \times 11)$$

is actually true. For another prime number $q_i > 11$, we have that

$$(1 + \frac{1}{q_i}) < (1 + \frac{1}{11})$$

and

$$\ln \ln(2^{19} \times 11) < \ln \ln(2^{19} \times q_i)$$

which clearly implies that the inequality (2.3) is true for every prime number $q_i \geq 11$. Now, suppose it is true for $m - 1$, with $m \geq 2$ and let us consider the assertion for those squarefree n with $\omega(n) = m$ [1]. So let $n = q_1 \times \cdots \times q_m$ be a squarefree number and assume that $q_1 < \cdots < q_m$ for $q_m \geq 11$.

Case 1: $q_m \geq \ln(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m) = \ln(2^{19} \times n)$.

By the induction hypothesis we have

$$\frac{\pi^2}{6} \times \frac{3}{2} \times (q_1 + 1) \times \cdots \times (q_{m-1} + 1) \leq e^\gamma \times q_1 \times \cdots \times q_{m-1} \times \ln \ln(2^{19} \times q_1 \times \cdots \times q_{m-1})$$

and hence

$$\frac{\pi^2}{6} \times \frac{3}{2} \times (q_1 + 1) \times \cdots \times (q_{m-1} + 1) \times (q_m + 1) \leq$$

$$e^\gamma \times q_1 \times \cdots \times q_{m-1} \times (q_m + 1) \times \ln \ln(2^{19} \times q_1 \times \cdots \times q_{m-1})$$

when we multiply the both sides of the inequality by $(q_m + 1)$. We want to show that

$$e^\gamma \times q_1 \times \cdots \times q_{m-1} \times (q_m + 1) \times \ln \ln(2^{19} \times q_1 \times \cdots \times q_{m-1}) \leq$$

$$e^\gamma \times q_1 \times \cdots \times q_{m-1} \times q_m \times \ln \ln(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m) = e^\gamma \times n \times \ln \ln(2^{19} \times n).$$

Indeed the previous inequality is equivalent with

$$q_m \times \ln \ln(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m) \geq (q_m + 1) \times \ln \ln(2^{19} \times q_1 \times \cdots \times q_{m-1})$$

or alternatively

$$\frac{q_m \times (\ln \ln(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m) - \ln \ln(2^{19} \times q_1 \times \cdots \times q_{m-1}))}{\ln q_m} \geq \frac{\ln \ln(2^{19} \times q_1 \times \cdots \times q_{m-1})}{\ln q_m}.$$

From the reference [1], we have that if $0 < a < b$, then

$$(2.4) \quad \frac{\ln b - \ln a}{b - a} = \frac{1}{(b - a)} \int_a^b \frac{dt}{t} > \frac{1}{b}.$$

We can apply the inequality (2.4) to the previous one just using $b = \ln(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m)$ and $a = \ln(2^{19} \times q_1 \times \cdots \times q_{m-1})$. Certainly, we have that

$$\begin{aligned} & \ln(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m) - \ln(2^{19} \times q_1 \times \cdots \times q_{m-1}) = \\ & \ln \frac{2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m}{2^{19} \times q_1 \times \cdots \times q_{m-1}} = \ln q_m. \end{aligned}$$

In this way, we obtain that

$$\frac{q_m \times (\ln \ln(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m) - \ln \ln(2^{19} \times q_1 \times \cdots \times q_{m-1}))}{\ln q_m} > \frac{q_m}{\ln(2^{19} \times q_1 \times \cdots \times q_m)}.$$

Using this result we infer that the original inequality is certainly satisfied if the next inequality is satisfied

$$\frac{q_m}{\ln(2^{19} \times q_1 \times \cdots \times q_m)} \geq \frac{\ln \ln(2^{19} \times q_1 \times \cdots \times q_{m-1})}{\ln q_m}$$

which is trivially true for $q_m \geq \ln(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m)$ [1].

Case 2: $q_m < \ln(2^{19} \times q_1 \times \cdots \times q_{m-1} \times q_m) = \ln(2^{19} \times n)$.

We need to prove that

$$\frac{\pi^2}{6} \times \frac{3}{2} \times \frac{\sigma(n)}{n} \leq e^\gamma \times \ln \ln(2^{19} \times n).$$

We know that $\frac{3}{2} < 1.503 < \frac{4}{2.66}$. Nevertheless, we could have that

$$\frac{3}{2} \times \frac{\sigma(n)}{n} \times \frac{\pi^2}{6} < \frac{4 \times \sigma(n)}{3 \times n} \times \frac{\pi^2}{2 \times 2.66}$$

and therefore, we only need to prove that

$$\frac{\sigma(3 \times n)}{3 \times n} \times \frac{\pi^2}{5.32} \leq e^\gamma \times \ln \ln(2^{19} \times n)$$

where this is possible because of $3 \nmid n$. If we apply the logarithm to the both sides of the inequality, then we obtain that

$$\ln\left(\frac{\pi^2}{5.32}\right) + (\ln(3+1) - \ln 3) + \sum_{j=1}^m (\ln(q_j + 1) - \ln q_j) \leq \gamma + \ln \ln \ln(2^{19} \times n).$$

From the reference [1], we note that

$$\ln(q_1 + 1) - \ln q_1 = \int_{q_1}^{q_1+1} \frac{dt}{t} < \frac{1}{q_1}.$$

In addition, note that $\ln\left(\frac{\pi^2}{5.32}\right) < \frac{1}{2} + 0.12$. However, we know that

$$\gamma + \ln \ln q_m < \gamma + \ln \ln \ln(2^{19} \times n)$$

since $q_m < \ln(2^{19} \times n)$ and therefore, it is enough to prove that

$$0.12 + \frac{1}{2} + \frac{1}{3} + \frac{1}{q_1} + \cdots + \frac{1}{q_m} \leq 0.12 + \sum_{q \leq q_m} \frac{1}{q} \leq \gamma + \ln \ln q_m$$

where $q_m \geq 11$. In this way, we only need to prove that

$$\sum_{q \leq q_m} \frac{1}{q} \leq \gamma + \ln \ln q_m - 0.12$$

which is true according to the Theorem 2.2 when $q_m \geq 11$. In this way, we finally show the Theorem is indeed satisfied. \square

Theorem 2.5. *Given a natural number*

$$n = 2^{a_1} \times 3^{a_2} \times 5^{a_3} \times 7^{a_4} > 5040$$

such that $a_1, a_2, a_3, a_4 \geq 0$ are integers, then the Robin's inequality is true for n .

Proof. Given a natural number $n = 2^{a_1} \times 3^{a_2} \times 5^{a_3} \times 7^{a_4} > 5040$ such that $a_1, a_2, a_3, a_4 \geq 0$ are integers, we need to prove that

$$\frac{\sigma(n)}{n} < e^\gamma \times \ln \ln n$$

that is true when

$$\prod_{i=1}^m \frac{q_i}{q_i - 1} < e^\gamma \times \ln \ln n$$

according to the inequality (2.1). Given a natural number $n = 2^{a_1} \times 3^{a_2} \times 5^{a_3} > 5040$ such that $a_1, a_2, a_3 \geq 0$ are integers, we have that

$$\prod_{i=1}^m \frac{q_i}{q_i - 1} \leq \frac{2 \times 3 \times 5}{1 \times 2 \times 4} = 3.75 < e^\gamma \times \ln \ln(5040) \approx 3.81.$$

However, we know for $n > 5040$ that

$$e^\gamma \times \ln \ln(5040) < e^\gamma \times \ln \ln n$$

and therefore, the proof is completed for that case. Hence, we only need to prove the Robin's inequality is true for every natural number $n = 2^{a_1} \times 3^{a_2} \times 5^{a_3} \times 7^{a_4} > 5040$ such that $a_1, a_2, a_3 \geq 0$ and $a_4 \geq 1$ are integers. In addition, we know the Robin's

inequality is true for every natural number $n > 5040$ such that $7^k \mid n$ and $7^7 \nmid n$ for some integer $1 \leq k \leq 6$ [3]. Therefore, we need to prove this case for those natural numbers $n > 5040$ such that $7^7 \mid n$. In this way, we have that

$$\prod_{i=1}^m \frac{q_i}{q_i - 1} \leq \frac{2 \times 3 \times 5 \times 7}{1 \times 2 \times 4 \times 6} = 4.375 < e^\gamma \times \ln \ln(7^7) \approx 4.65.$$

However, we know for $n > 5040$ and $7^7 \mid n$ that

$$e^\gamma \times \ln \ln(7^7) \leq e^\gamma \times \ln \ln n$$

and as a consequence, the proof is completed. \square

Theorem 2.6. *The Robin's inequality is true for every natural number $n > 5040$ when $3 \nmid n$. More precisely: every possible counterexample $n > 5040$ of the Robin's inequality must comply that $(2^{20} \times 3^{13}) \mid n$.*

Proof. We will check the Robin's inequality is true for every natural number $n = q_1^{a_1} \times q_2^{a_2} \times \cdots \times q_m^{a_m} > 5040$ such that q_1, q_2, \dots, q_m are prime numbers, a_1, a_2, \dots, a_m are positive integers and $3 \nmid n$. We know this is true when the greatest prime divisor of $n > 5040$ is lesser than or equal to 7 according to the Theorem 2.5. Therefore, the remaining case is when the greatest prime divisor of $n > 5040$ is greater than 7. We need to prove that

$$\frac{\sigma(n)}{n} < e^\gamma \times \ln \ln n$$

that is true when

$$\frac{\pi^2}{6} \times \prod_{i=1}^m \frac{q_i + 1}{q_i} \leq e^\gamma \times \ln \ln n$$

according to the Theorem 2.1. Using the equation (2.2), we obtain that will be equivalent to

$$\frac{\pi^2}{6} \times \frac{\sigma(n')}{n'} < e^\gamma \times \ln \ln n$$

where $n' = q_1 \times \cdots \times q_m$ is the squarefree kernel of n [1]. However, the Robin's inequality has been proved for all integers n not divisible by 2 (which are bigger than 10) [1]. Hence, we only need to prove the Robin's inequality is true when $2 \mid n'$. In addition, we know the Robin's inequality is true for every natural number $n > 5040$ such that $2^k \mid n$ and $2^{20} \nmid n$ for some integer $1 \leq k \leq 19$ [3]. Consequently, we only need to prove the Robin's inequality is true for all $n > 5040$ such that $2^{20} \mid n$ and thus,

$$e^\gamma \times n' \times \ln \ln(2^{19} \times \frac{n'}{2}) < e^\gamma \times n' \times \ln \ln n$$

because of $2^{19} \times \frac{n'}{2} < n$ when $2^{20} \mid n$ and $2 \mid n'$. In this way, we only need to prove that

$$\frac{\pi^2}{6} \times \sigma(n') \leq e^\gamma \times n' \times \ln \ln(2^{19} \times \frac{n'}{2}).$$

According to the equation (2.2) and $2 \mid n'$, we have that

$$\frac{\pi^2}{6} \times 3 \times \sigma(\frac{n'}{2}) \leq e^\gamma \times 2 \times \frac{n'}{2} \times \ln \ln(2^{19} \times \frac{n'}{2})$$

which is the same as

$$\frac{\pi^2}{6} \times \frac{3}{2} \times \sigma(\frac{n'}{2}) \leq e^\gamma \times \frac{n'}{2} \times \ln \ln(2^{19} \times \frac{n'}{2})$$

that is true according to the Theorem 2.4 when $3 \nmid \frac{n'}{2}$. In addition, we know the Robin's inequality is true for every natural number $n > 5040$ such that $3^k \mid n$ and $3^{13} \nmid n$ for some integer $1 \leq k \leq 12$ [3]. Consequently, we only need to prove the Robin's inequality is true for all $n > 5040$ such that $2^{20} \mid n$ and $3^{13} \mid n$. To sum up, the proof is completed. \square

Theorem 2.7. *The Robin's inequality is true for every natural number $n > 5040$ when n is not divisible by 5. Moreover, we show the Robin's inequality is true for every natural number $n > 5040$ when n is not divisible by 7.*

Proof. Let's define $s(n) = \frac{\sigma(n)}{n}$ [5]. Hence, we need to prove that

$$s(n) < e^\gamma \times \ln \ln n$$

when $(2^{20} \times 3^{13}) \mid n$. Suppose that $n = 2^a \times 3^b \times m$, where $a \geq 20$, $b \geq 13$, $2 \nmid m$, $3 \nmid m$ and $5 \nmid m$ or $7 \nmid m$. Therefore, we need to prove that

$$s(2^a \times 3^b \times m) < e^\gamma \times \ln \ln(2^a \times 3^b \times m).$$

We know that

$$s(2^a \times 3^b \times m) = s(3^b) \times s(2^a \times m)$$

since s is multiplicative [5]. In addition, we know that $s(3^b) < \frac{3}{2}$ for every positive integer b [5]. In this way, we have that

$$s(3^b) \times s(2^a \times m) < \frac{3}{2} \times s(2^a \times m).$$

Now, consider that

$$\frac{3}{2} \times s(2^a \times m) = \frac{9}{8} \times s(3) \times s(2^a \times m) = \frac{9}{8} \times s(2^a \times 3 \times m)$$

where $s(3) = \frac{4}{3}$ since s is multiplicative [5]. Nevertheless, we have that

$$\frac{9}{8} \times s(2^a \times 3 \times m) < s(5) \times s(2^a \times 3 \times m) = s(2^a \times 3 \times 5 \times m)$$

and

$$\frac{9}{8} \times s(2^a \times 3 \times m) < s(7) \times s(2^a \times 3 \times m) = s(2^a \times 3 \times 7 \times m)$$

where $5 \nmid m$ or $7 \nmid m$, $s(5) = \frac{6}{5}$ and $s(7) = \frac{8}{7}$. However, we know the Robin's inequality is true for $2^a \times 3 \times 5 \times m$ and $2^a \times 3 \times 7 \times m$ when $a \geq 20$, since this is true for every natural number $n > 5040$ such that $3^k \mid n$ and $3^{13} \nmid n$ for some integer $1 \leq k \leq 12$ [3]. Hence, we would have that

$$s(2^a \times 3 \times 5 \times m) < e^\gamma \times \ln \ln(2^a \times 3 \times 5 \times m) < e^\gamma \times \ln \ln(2^a \times 3^b \times m)$$

and

$$s(2^a \times 3 \times 7 \times m) < e^\gamma \times \ln \ln(2^a \times 3 \times 7 \times m) < e^\gamma \times \ln \ln(2^a \times 3^b \times m)$$

when $b \geq 13$. \square

Theorem 2.8. *The Robin's inequality is true for every natural number $n > 5040$ when n is not divisible by any prime number $11 \leq q_m \leq 47$.*

Proof. We need to prove that

$$s(n) < e^\gamma \times \ln \ln n$$

when $(2^{20} \times 3^{13} \times 7^7) \mid n$. Suppose that $n = 2^a \times 3^b \times 7^c \times m$, where $a \geq 20$, $b \geq 13$, $c \geq 7$, $2 \nmid m$, $3 \nmid m$, $7 \nmid m$, $q_m \nmid m$ and $11 \leq q_m \leq 47$. Therefore, we need to prove that

$$s(2^a \times 3^b \times 7^c \times m) < e^\gamma \times \ln \ln(2^a \times 3^b \times 7^c \times m).$$

We know that

$$s(2^a \times 3^b \times 7^c \times m) = s(7^c) \times s(2^a \times 3^b \times m)$$

since s is multiplicative [5]. In addition, we know that $s(7^c) < \frac{7}{6}$ for every positive integer c [5]. In this way, we have that

$$s(7^c) \times s(2^a \times 3^b \times m) < \frac{7}{6} \times s(2^a \times 3^b \times m).$$

However, that would be equivalent to

$$\frac{49}{48} \times s(7) \times s(2^a \times 3^b \times m) = \frac{49}{48} \times s(2^a \times 3^b \times 7 \times m)$$

where $s(7) = \frac{8}{7}$. In addition, we know that

$$\frac{49}{48} \times s(2^a \times 3^b \times 7 \times m) < s(q_m) \times s(2^a \times 3^b \times 7 \times m) = s(2^a \times 3^b \times 7 \times q_m \times m)$$

where $q_m \nmid m$, $s(q_m) = \frac{q_m+1}{q_m}$ and $11 \leq q_m \leq 47$. Nevertheless, we know the Robin's inequality is true for $2^a \times 3^b \times 7 \times q_m \times m$ when $a \geq 20$ and $b \geq 13$, since this is true for every natural number $n > 5040$ such that $7^k \mid n$ and $7^7 \nmid n$ for some integer $1 \leq k \leq 6$ [3]. Hence, we would have that

$$s(2^a \times 3^b \times 7 \times q_m \times m) < e^\gamma \times \ln \ln(2^a \times 3^b \times 7 \times q_m \times m) < e^\gamma \times \ln \ln(2^a \times 3^b \times 7^c \times m)$$

when $c \geq 7$ and $11 \leq q_m \leq 47$. \square

Theorem 2.9. *The Robin's inequality is true for every natural number $n > 5040$ when n is not divisible by any prime number $53 \leq q_m \leq 113$.*

Proof. We know the Robin's inequality is true for every natural number $n > 5040$ such that $11^k \mid n$ and $11^6 \nmid n$ for some integer $1 \leq k \leq 5$ [3]. We need to prove that

$$s(n) < e^\gamma \times \ln \ln n$$

when $(2^{20} \times 3^{13} \times 11^6) \mid n$. Suppose that $n = 2^a \times 3^b \times 11^c \times m$, where $a \geq 20$, $b \geq 13$, $c \geq 6$, $2 \nmid m$, $3 \nmid m$, $11 \nmid m$, $q_m \nmid m$ and $53 \leq q_m \leq 113$. Therefore, we need to prove that

$$s(2^a \times 3^b \times 11^c \times m) < e^\gamma \times \ln \ln(2^a \times 3^b \times 11^c \times m).$$

We know that

$$s(2^a \times 3^b \times 11^c \times m) = s(11^c) \times s(2^a \times 3^b \times m)$$

since s is multiplicative [5]. In addition, we know that $s(11^c) < \frac{11}{10}$ for every positive integer c [5]. In this way, we have that

$$s(11^c) \times s(2^a \times 3^b \times m) < \frac{11}{10} \times s(2^a \times 3^b \times m).$$

However, that would be equivalent to

$$\frac{121}{120} \times s(11) \times s(2^a \times 3^b \times m) = \frac{121}{120} \times s(2^a \times 3^b \times 11 \times m)$$

where $s(11) = \frac{12}{11}$. In addition, we know that

$$\frac{121}{120} \times s(2^a \times 3^b \times 11 \times m) < s(q_m) \times s(2^a \times 3^b \times 11 \times m) = s(2^a \times 3^b \times 11 \times q_m \times m)$$

where $q_m \nmid m$, $s(q_m) = \frac{q_m+1}{q_m}$ and $53 \leq q_m \leq 113$. Nevertheless, we know the Robin's inequality is true for $2^a \times 3^b \times 11 \times q_m \times m$ when $a \geq 20$ and $b \geq 13$, since this is true for every natural number $n > 5040$ such that $11^k \mid n$ and $11^6 \nmid n$ for some integer $1 \leq k \leq 5$ [3]. Hence, we would have that

$$s(2^a \times 3^b \times 11 \times q_m \times m) < e^\gamma \times \ln \ln(2^a \times 3^b \times 11 \times q_m \times m) < e^\gamma \times \ln \ln(2^a \times 3^b \times 11^c \times m)$$

when $c \geq 6$ and $53 \leq q_m \leq 113$. \square

Theorem 2.10. *The Robin's inequality is true for every natural number $n > 5040$ when n is not divisible by any prime number $q_m \leq 113$.*

Proof. This is a compendium of the results from the Theorems 2.6, 2.7, 2.8 and 2.9. \square

Theorem 2.11. *The Robin's inequality is true for every natural number $n = 113^k \times n' > 5040$ when $(\ln n')^\beta \leq \ln n$ such that $\beta = \frac{113}{112}$, $113 \nmid n'$ and for an integer $k \geq 1$.*

Proof. Suppose that $n = 113^k \times n'$ for an integer $k \geq 1$ and a natural number n' , where $113 \nmid n'$. Hence, we need to prove that

$$(2.5) \quad s(n) < e^\gamma \times \ln \ln n.$$

Now, consider the case when $n' \leq 5040$. We know that

$$(2.6) \quad s(113^k \times n') = s(113^k) \times s(n')$$

since s is multiplicative [5]. In addition, we know that $s(113^k) < \frac{113}{112}$ for every possible integer $k \geq 1$ [5]. In this way, we have that

$$(2.7) \quad s(113^k) \times s(n') < \frac{113}{112} \times s(n').$$

However, we know the maximum value of $s(n')$ is when $n' = 5040$, where $n' \leq 5040$ and $s(5040)$ has the following property of $s(5040) < 3.9$. Consequently, we have that

$$\frac{113}{112} \times s(n') < \frac{113}{112} \times 3.9$$

when $n' \leq 5040$. We know that $\frac{113}{112} \times 3.9 < 4 < e^\gamma \times \ln \ln e^{e^4}$. In addition, we know the Robin's inequality is true for every natural number $5040 < n \leq 10^{10^{10}}$ [3]. Therefore, we only need to prove this case for every natural number $n > 10^{10^{10}}$. Nevertheless, for every natural number $n > 10^{10^{10}}$, we will have that

$$e^\gamma \times \ln \ln e^{e^4} < e^\gamma \times \ln \ln n$$

due to $e^{e^4} < 10^{10^{10}}$ and as result, the Theorem is true for this case.

In the case of $n' > 5040$, we know that

$$s(n') < e^\gamma \times \ln \ln n'$$

because of $113 \nmid n'$ after of applying the result of Theorem 2.10. If we multiply the both sides of the inequality by $\frac{113}{112}$, then we would have that

$$\frac{113}{112} \times s(n') < \frac{113}{112} \times e^\gamma \times \ln \ln n'.$$

If we use the inequalities (2.6) and (2.7), then we obtain that

$$s(n) < \frac{113}{112} \times e^\gamma \times \ln \ln n'.$$

and thus, the inequality (2.5) is true when

$$\frac{113}{112} \times \ln \ln n' \leq \ln \ln n$$

is true, that is equivalent to

$$(\ln n')^{\frac{113}{112}} \leq \ln n.$$

□

Theorem 2.12. *Given a natural number*

$$n = q_1^{a_1} \times q_2^{a_2} \times \cdots \times q_m^{a_m}$$

such that $n > 5040$, q_1, q_2, \dots, q_m are prime numbers and a_1, a_2, \dots, a_m are positive integers, then the Robin's inequality is true for n when $q_1^\alpha \times q_2^\alpha \times \cdots \times q_m^\alpha \leq n$, where $\alpha = (\ln n')^\beta$, $\beta = (\frac{\pi^2}{6} - 1)$ and n' is the squarefree kernel of n .

Proof. We will check the Robin's inequality for every natural number $n = q_1^{a_1} \times q_2^{a_2} \times \cdots \times q_m^{a_m} > 5040$ such that q_1, q_2, \dots, q_m are prime numbers and a_1, a_2, \dots, a_m are positive integers. We need to prove that

$$\frac{\sigma(n)}{n} < e^\gamma \times \ln \ln n$$

that is true when

$$\frac{\pi^2}{6} \times \prod_{i=1}^m \frac{q_i + 1}{q_i} \leq e^\gamma \times \ln \ln n$$

according to the Theorem 2.1. Using the equation (2.2), we obtain that will be equivalent to

$$\frac{\pi^2}{6} \times \frac{\sigma(n')}{n'} \leq e^\gamma \times \ln \ln n$$

where $n' = q_1 \times \cdots \times q_m$ is the squarefree kernel of n [1]. However, the Robin's inequality has been proved for all the squarefree integers $n' \notin \{2, 3, 5, 6, 10, 30\}$ [1]. In addition, as a consequence of the Theorem 2.5, the Robin's inequality is true for every natural number $n > 5040$ when $n' \in \{2, 3, 5, 6, 10, 30\}$ such that n' is the squarefree kernel of n . In this way, we have that

$$\frac{\sigma(n')}{n'} < e^\gamma \times \ln \ln n'$$

and therefore, it is enough to prove that

$$\frac{\pi^2}{6} \times e^\gamma \times \ln \ln n' \leq e^\gamma \times \ln \ln n$$

which is simplified as

$$\frac{\pi^2}{6} \times \ln \ln n' \leq \ln \ln n$$

and

$$\ln(\ln n')^{\frac{\pi^2}{6}} \leq \ln \ln n$$

that is true when

$$(\ln n')^{\frac{\pi^2}{6}} \leq \ln n$$

is true. Consequently, that would be equivalent to

$$e^{(\ln n')^{\frac{\pi^2}{6}}} \leq n$$

which is equal to

$$e^{(\ln n') \times (\ln n')^{\left(\frac{\pi^2}{6}-1\right)}} \leq n$$

that is

$$n'^{(\ln n')^{\left(\frac{\pi^2}{6}-1\right)}} \leq n$$

and therefore, the proof is completed. \square

3. CONCLUSIONS

The practical uses of the Riemann hypothesis include many propositions which are known true under the Riemann hypothesis, and some that can be shown equivalent to the Riemann hypothesis [2]. Certainly, the Riemann hypothesis is close related to various mathematical topics such as the distribution of prime numbers, the growth of arithmetic functions, the Lindelöf hypothesis, the large prime gap conjecture, etc [2]. Indeed, a proof of the Riemann hypothesis could spur considerable advances in many mathematical areas, such as the number theory and pure mathematics [2]. In this way, we made a new step forward in the efforts of trying to prove the Riemann hypothesis.

REFERENCES

- [1] YoungJu Choie, Nicolas Lichiardopol, Pieter Moree, and Patrick Solé. On Robin's criterion for the Riemann hypothesis. *Journal de Théorie des Nombres de Bordeaux*, 19(2):357–372, 2007.
- [2] Keith J. Devlin. *The Millennium Problems: The Seven Greatest Unsolved Mathematical Puzzles Of Our Time*. Basic Books, 2003.
- [3] Alexander Hertlein. Robin's Inequality for New Families of Integers. *Integers*, 18, 2018.
- [4] J. Barkley Rosser and Lowell Schoenfeld. Approximate Formulas for Some Functions of Prime Numbers. *Illinois Journal of Mathematics*, 6(1):64–94, 1962.
- [5] Robert Vojak. On numbers satisfying Robin's inequality, properties of the next counterexample and improved specific bounds. *arXiv preprint arXiv:2005.09307*, 2020.
- [6] David G. Wells. *Prime Numbers, The Most Mysterious Figures in Math*. John Wiley & Sons, Inc., 2005.

COPSONIC, 1471 ROUTE DE SAINT-NAUPHARY 82000 MONTAUBAN, FRANCE
E-mail address: vega.frank@gmail.com